

# Quark and Lepton Compositeness, Searches for

The latest unpublished results are described in the “Quark and Lepton Compositeness” review.

A REVIEW GOES HERE – Check our WWW List of Reviews

---

## CONTENTS:

- Scale Limits for Contact Interactions:  $\Lambda(eeee)$
  - Scale Limits for Contact Interactions:  $\Lambda(ee\mu\mu)$
  - Scale Limits for Contact Interactions:  $\Lambda(ee\tau\tau)$
  - Scale Limits for Contact Interactions:  $\Lambda(\ell\ell\ell\ell)$
  - Scale Limits for Contact Interactions:  $\Lambda(eeqq)$
  - Scale Limits for Contact Interactions:  $\Lambda(\mu\mu qq)$
  - Scale Limits for Contact Interactions:  $\Lambda(\ell\nu\ell\nu)$
  - Scale Limits for Contact Interactions:  $\Lambda(e\nu qq)$
  - Scale Limits for Contact Interactions:  $\Lambda(qqqq)$
  - Scale Limits for Contact Interactions:  $\Lambda(\nu\nu qq)$
  - Mass Limits for Excited  $e$  ( $e^*$ )
    - Limits for Excited  $e$  ( $e^*$ ) from Pair Production
    - Limits for Excited  $e$  ( $e^*$ ) from Single Production
    - Limits for Excited  $e$  ( $e^*$ ) from  $e^+e^- \rightarrow \gamma\gamma$
    - Indirect Limits for Excited  $e$  ( $e^*$ )
  - Mass Limits for Excited  $\mu$  ( $\mu^*$ )
    - Limits for Excited  $\mu$  ( $\mu^*$ ) from Pair Production
    - Limits for Excited  $\mu$  ( $\mu^*$ ) from Single Production
    - Indirect Limits for Excited  $\mu$  ( $\mu^*$ )
  - Mass Limits for Excited  $\tau$  ( $\tau^*$ )
    - Limits for Excited  $\tau$  ( $\tau^*$ ) from Pair Production
    - Limits for Excited  $\tau$  ( $\tau^*$ ) from Single Production
  - Mass Limits for Excited Neutrino ( $\nu^*$ )
    - Limits for Excited  $\nu$  ( $\nu^*$ ) from Pair Production
    - Limits for Excited  $\nu$  ( $\nu^*$ ) from Single Production
  - Mass Limits for Excited  $q$  ( $q^*$ )
    - Limits for Excited  $q$  ( $q^*$ ) from Pair Production
    - Limits for Excited  $q$  ( $q^*$ ) from Single Production
  - Mass Limits for Color Sextet Quarks ( $q_6$ )
  - Mass Limits for Color Octet Charged Leptons ( $\ell_8$ )
  - Mass Limits for Color Octet Neutrinos ( $\nu_8$ )
  - Mass Limits for  $W_8$  (Color Octet  $W$  Boson)
- 

## SCALE LIMITS for Contact Interactions: $\Lambda(eeee)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;8.3</b>	<b>&gt;10.3</b>	95	<sup>1</sup> BOURILKOV	01	RVUE $E_{\text{cm}} = 192\text{--}208$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

>4.5	>7.0	95	<sup>2</sup> SCHAEL	07A	ALEP	$E_{\text{cm}} = 189\text{--}209$ GeV
>5.3	>6.8	95	ABDALLAH	06C	DLPH	$E_{\text{cm}} = 130\text{--}207$ GeV
>4.7	>6.1	95	<sup>3</sup> ABBIENDI	04G	OPAL	$E_{\text{cm}} = 130\text{--}207$ GeV
>4.4	>5.4	95	ABREU	00S	DLPH	$E_{\text{cm}} = 183\text{--}189$ GeV
>4.3	>4.9	95	ACCIARRI	00P	L3	$E_{\text{cm}} = 130\text{--}189$ GeV

<sup>1</sup> A combined analysis of the data from ALEPH, DELPHI, L3, and OPAL.

<sup>2</sup> SCHAEL 07A limits are from  $R_c$ ,  $Q_{FB}^{depl}$ , and hadronic cross section measurements.

<sup>3</sup> ABBIENDI 04G limits are from  $e^+e^- \rightarrow e^+e^-$  cross section at  $\sqrt{s} = 130\text{--}207$  GeV.

### SCALE LIMITS for Contact Interactions: $\Lambda(ee\mu\mu)$

Limits are for  $\Lambda_{LL}^{\pm}$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>6.6	<b>&gt;9.5</b>	95	<sup>4</sup> SCHAEL	07A	ALEP $E_{\text{cm}} = 189\text{--}209$ GeV
<b>&gt; 8.5</b>	>3.8	95	ACCIARRI	00P	L3 $E_{\text{cm}} = 130\text{--}189$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

>7.3	>7.6	95	ABDALLAH	06C	DLPH $E_{\text{cm}} = 130\text{--}207$ GeV
>8.1	>7.3	95	<sup>5</sup> ABBIENDI	04G	OPAL $E_{\text{cm}} = 130\text{--}207$ GeV
>6.6	>6.3	95	ABREU	00S	DLPH $E_{\text{cm}} = 183\text{--}189$ GeV

<sup>4</sup> SCHAEL 07A limits are from  $R_c$ ,  $Q_{FB}^{depl}$ , and hadronic cross section measurements.

<sup>5</sup> ABBIENDI 04G limits are from  $e^+e^- \rightarrow \mu\mu$  cross section at  $\sqrt{s} = 130\text{--}207$  GeV.

### SCALE LIMITS for Contact Interactions: $\Lambda(ee\tau\tau)$

Limits are for  $\Lambda_{LL}^{\pm}$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;7.9</b>	>5.8	95	<sup>6</sup> SCHAEL	07A	ALEP $E_{\text{cm}} = 189\text{--}209$ GeV
<b>&gt;7.9</b>	>4.6	95	ABDALLAH	06C	DLPH $E_{\text{cm}} = 130\text{--}207$ GeV
>4.9	<b>&gt;7.2</b>	95	<sup>7</sup> ABBIENDI	04G	OPAL $E_{\text{cm}} = 130\text{--}207$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

>5.2	>5.4	95	ABREU	00S	DLPH $E_{\text{cm}} = 183\text{--}189$ GeV
>5.4	>4.7	95	ACCIARRI	00P	L3 $E_{\text{cm}} = 130\text{--}189$ GeV

<sup>6</sup> SCHAEL 07A limits are from  $R_c$ ,  $Q_{FB}^{depl}$ , and hadronic cross section measurements.

<sup>7</sup> ABBIENDI 04G limits are from  $e^+e^- \rightarrow \tau\tau$  cross section at  $\sqrt{s} = 130\text{--}207$  GeV.

### SCALE LIMITS for Contact Interactions: $\Lambda(llll)$

Lepton universality assumed. Limits are for  $\Lambda_{LL}^{\pm}$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>7.9	<b>&gt; 10.3</b>	95	<sup>8</sup> SCHAEL	07A	ALEP $E_{\text{cm}} = 189\text{--}209$ GeV
<b>&gt;9.1</b>	>8.2	95	ABDALLAH	06C	DLPH $E_{\text{cm}} = 130\text{--}207$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

>7.7	>9.5	95	<sup>9</sup> ABBIENDI	04G	OPAL	$E_{\text{cm}} = 130\text{--}207$ GeV
			<sup>10</sup> BABICH	03	RVUE	
>9.0	>5.2	95	ACCIARRI	00P	L3	$E_{\text{cm}} = 130\text{--}189$ GeV

<sup>8</sup> SCHAEL 07A limits are from  $R_c$ ,  $Q_{FB}^{depl}$ , and hadronic cross section measurements.

<sup>9</sup> ABBIENDI 04G limits are from  $e^+e^- \rightarrow \ell^+\ell^-$  cross section at  $\sqrt{s} = 130\text{--}207$  GeV.

<sup>10</sup> BABICH 03 obtain a bound  $-0.175 \text{ TeV}^{-2} < 1/\Lambda_{LL}^2 < 0.095 \text{ TeV}^{-2}$  (95%CL) in a model independent analysis allowing all of  $\Lambda_{LL}$ ,  $\Lambda_{LR}$ ,  $\Lambda_{RL}$ ,  $\Lambda_{RR}$  to coexist.

### SCALE LIMITS for Contact Interactions: $\Lambda(eeqq)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 8.4	<b>&gt;10.2</b>	95	<sup>11</sup> ABDALLAH	09	DLPH ( <i>eebb</i> )
<b>&gt; 9.4</b>	<b>&gt;5.6</b>	95	<sup>12</sup> SCHAEL	07A	ALEP ( <i>eecc</i> )
> 9.4	>4.9	95	<sup>11</sup> SCHAEL	07A	ALEP ( <i>eebb</i> )
<b>&gt;23.3</b>	<b>&gt;12.5</b>	95	<sup>13</sup> CHEUNG	01B	RVUE ( <i>eeuu</i> )
<b>&gt;11.1</b>	<b>&gt;26.4</b>	95	<sup>13</sup> CHEUNG	01B	RVUE ( <i>eedd</i> )

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 4.2	>4.0	95	<sup>14</sup> AARON	11C	H1 ( <i>eeqq</i> )
> 3.8	>3.8	95	<sup>15</sup> ABDALLAH	11	DLPH ( <i>eecc</i> )
>12.9	>7.2	95	<sup>16</sup> SCHAEL	07A	ALEP ( <i>eeqq</i> )
> 3.7	>5.9	95	<sup>17</sup> ABULENCIA	06L	CDF ( <i>eeqq</i> )

<sup>11</sup> ABDALLAH 09 and SCHAEL 07A limits are from  $R_b$ ,  $A_{FB}^b$ .

<sup>12</sup> SCHAEL 07A limits are from  $R_c$ ,  $Q_{FB}^{depl}$ , and hadronic cross section measurements.

<sup>13</sup> CHEUNG 01B is an update of BARGER 98E.

<sup>14</sup> AARON 11C limits are from  $Q^2$  spectrum measurements of  $e^\pm p \rightarrow e^\pm X$ .

<sup>15</sup> ABDALLAH 11 limit is from  $e^+e^- \rightarrow t\bar{c}$  cross section.  $\Lambda_{LL} = \Lambda_{LR} = \Lambda_{RL} = \Lambda_{RR}$  is assumed.

<sup>16</sup> SCHAEL 07A limit assumes quark flavor universality of the contact interactions.

<sup>17</sup> ABULENCIA 06L limits are from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV.

### SCALE LIMITS for Contact Interactions: $\Lambda(\mu\mu qq)$

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;4.5</b>	<b>&gt;4.9</b>	95	<sup>18</sup> AAD	11E	ATLS ( $\mu\mu qq$ ) (isosinglet)

• • • We do not use the following data for averages, fits, limits, etc. • • •

>2.9	>4.2	95	<sup>19</sup> ABE	97T	CDF ( $\mu\mu qq$ ) (isosinglet)
------	------	----	-------------------	-----	----------------------------------

<sup>18</sup> AAD 11E limits are from  $\mu^+\mu^-$  mass distribution in  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV.

<sup>19</sup> ABE 97T limits are from  $\mu^+\mu^-$  mass distribution in  $\bar{p}p \rightarrow \mu^+\mu^- X$  at  $E_{\text{cm}} = 1.8$  TeV.

### SCALE LIMITS for Contact Interactions: $\Lambda(\ell\nu\ell\nu)$

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;3.10</b>	90	<sup>20</sup> JODIDIO	86	SPEC $\Lambda_{LR}^\pm(\nu_\mu\nu_e\mu e)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>3.8	21 DIAZCRUZ	94	RVUE	$\Lambda_{LL}^+(\tau\nu_\tau e\nu_e)$
>8.1	21 DIAZCRUZ	94	RVUE	$\Lambda_{LL}^-(\tau\nu_\tau e\nu_e)$
>4.1	22 DIAZCRUZ	94	RVUE	$\Lambda_{LL}^+(\tau\nu_\tau \mu\nu_\mu)$
>6.5	22 DIAZCRUZ	94	RVUE	$\Lambda_{LL}^-(\tau\nu_\tau \mu\nu_\mu)$

<sup>20</sup> JODIDIO 86 limit is from  $\mu^+ \rightarrow \bar{\nu}_\mu e^+ \nu_e$ . Chirality invariant interactions  $L = (g^2/\Lambda^2) [\eta_{LL} (\bar{\nu}_\mu L \gamma^\alpha \mu_L) (\bar{e}_L \gamma_\alpha \nu_{eL}) + \eta_{LR} (\bar{\nu}_\mu L \gamma^\alpha \nu_{eL}) (\bar{e}_R \gamma_\alpha \mu_R)]$  with  $g^2/4\pi = 1$  and  $(\eta_{LL}, \eta_{LR}) = (0, \pm 1)$  are taken. No limits are given for  $\Lambda_{LL}^\pm$  with  $(\eta_{LL}, \eta_{LR}) = (\pm 1, 0)$ . For more general constraints with right-handed neutrinos and chirality nonconserving contact interactions, see their text.

<sup>21</sup> DIAZCRUZ 94 limits are from  $\Gamma(\tau \rightarrow e\nu\nu)$  and assume flavor-dependent contact interactions with  $\Lambda(\tau\nu_\tau e\nu_e) \ll \Lambda(\mu\nu_\mu e\nu_e)$ .

<sup>22</sup> DIAZCRUZ 94 limits are from  $\Gamma(\tau \rightarrow \mu\nu\nu)$  and assume flavor-dependent contact interactions with  $\Lambda(\tau\nu_\tau \mu\nu_\mu) \ll \Lambda(\mu\nu_\mu e\nu_e)$ .

### SCALE LIMITS for Contact Interactions: $\Lambda(e\nu q q)$

VALUE (TeV)	CL%	DOCUMENT ID	TECN
<b>&gt;2.81</b>	95	<sup>23</sup> AFFOLDER 01I	CDF

<sup>23</sup> AFFOLDER 00I bound is for a scalar interaction  $\bar{q}_R q_L \bar{\nu} e_L$ .

### SCALE LIMITS for Contact Interactions: $\Lambda(qqqq)$

Limits are for  $\Lambda_{LL}^\pm$  with color-singlet isoscalar exchanges among  $u_L$ 's and  $d_L$ 's only, unless otherwise noted. See EICHTEN 84 for details.

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;5.6</b>	95	<sup>24</sup> KHACHATRY...11F	CMS	$pp \rightarrow$ dijet angl.; $\Lambda_{LL}^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>3.4	95	<sup>25</sup> AAD	11	ATLS	$pp \rightarrow$ dijet; $\Lambda_{LL}^+$
>4.0	95	<sup>26</sup> KHACHATRY...10A	CMS		$pp$ ; dijet centrality; $\Lambda_{LL}^+$
>2.96	95	<sup>27</sup> ABAZOV	09AE	D0	$p\bar{p} \rightarrow$ dijet, angl. $\Lambda_{LL}^+$

<sup>24</sup> KHACHATRYAN 11F limit is from dijet angular distribution in  $pp$  collisions at  $E_{cm} = 7$  TeV. They also obtain  $\Lambda_{LL}^- > 6.7$  TeV.

<sup>25</sup> AAD 11 limit is from dijet angular distribution and dijet centrality ratio in  $pp$  collisions at  $E_{cm} = 7$  TeV.

<sup>26</sup> The quoted limit is from dijet centrality ratio measurement in  $pp$  collisions at  $\sqrt{s}=7$  TeV.

<sup>27</sup> ABAZOV 09AE also obtain  $\Lambda_{LL}^- > 2.96$  TeV.

### SCALE LIMITS for Contact Interactions: $\Lambda(\nu\nu q q)$

Limits are for  $\Lambda_{LL}^\pm$  only. For other cases, see each reference.

$\Lambda_{LL}^+$ (TeV)	$\Lambda_{LL}^-$ (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;5.0</b>	<b>&gt;5.4</b>	95	<sup>28</sup> MCFARLAND 98	CCFR	$\nu N$ scattering

<sup>28</sup> MCFARLAND 98 assumed a flavor universal interaction. Neutrinos were mostly of muon type.

## MASS LIMITS for Excited $e$ ( $e^*$ )

Most  $e^+e^-$  experiments assume one-photon or  $Z$  exchange. The limits from some  $e^+e^-$  experiments which depend on  $\lambda$  have assumed transition couplings which are chirality violating ( $\eta_L = \eta_R$ ). However they can be interpreted as limits for chirality-conserving interactions after multiplying the coupling value  $\lambda$  by  $\sqrt{2}$ ; see Note.

Excited leptons have the same quantum numbers as other ortholeptons. See also the searches for ortholeptons in the "Searches for Heavy Leptons" section.

### Limits for Excited $e$ ( $e^*$ ) from Pair Production

These limits are obtained from  $e^+e^- \rightarrow e^{*+}e^{*-}$  and thus rely only on the (electroweak) charge of  $e^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $e^*$  coupling is assumed to be of sequential type. Possible  $t$  channel contribution from transition magnetic coupling is neglected. All limits assume a dominant  $e^* \rightarrow e\gamma$  decay except the limits from  $\Gamma(Z)$ .

For limits prior to 1987, see our 1992 edition (Physical Review **D45** S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;103.2</b>	95	<sup>29</sup> ABBIENDI	02G OPAL	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>102.8	95	<sup>30</sup> ACHARD	03B L3	$e^+e^- \rightarrow e^*e^*$ Homodoublet type
<sup>29</sup> From $e^+e^-$ collisions at $\sqrt{s} = 183\text{--}209$ GeV. $f = f'$ is assumed.				
<sup>30</sup> From $e^+e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = f'$ is assumed. ACHARD 03B also obtain limit for $f = -f'$ : $m_{e^*} > 96.6$ GeV.				

### Limits for Excited $e$ ( $e^*$ ) from Single Production

These limits are from  $e^+e^- \rightarrow e^*e$ ,  $W \rightarrow e^*\nu$ , or  $ep \rightarrow e^*X$  and depend on transition magnetic coupling between  $e$  and  $e^*$ . All limits assume  $e^* \rightarrow e\gamma$  decay except as noted. Limits from LEP, UA2, and H1 are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda\text{--}m_{e^*}$  plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45** S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;1070</b>	95	<sup>31</sup> CHATRCHYAN	11X CMS	$pp \rightarrow ee^*X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 272	95	<sup>32</sup> AARON	08A H1	$ep \rightarrow e^*X$
		<sup>33</sup> ABAZOV	08H D0	$p\bar{p} \rightarrow e^*e$
> 209	95	<sup>34</sup> ACOSTA	05B CDF	$p\bar{p} \rightarrow e^*X$
> 206	95	<sup>35</sup> ACHARD	03B L3	$e^+e^- \rightarrow ee^*$
> 208	95	<sup>36</sup> ABBIENDI	02G OPAL	$e^+e^- \rightarrow ee^*$
> 228	95	<sup>37</sup> CHEKANOV	02D ZEUS	$ep \rightarrow e^*X$

- 31 CHATRCHYAN 11X search for single  $e^*$  production in  $pp$  collisions with the decay  $e^* \rightarrow e\gamma$ .  $f = f' = \Lambda/m_{e^*}$  is assumed. See their Fig. 2 for the exclusion plot in the mass-coupling plane.
- 32 AARON 08A search for single  $e^*$  production in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma$ ,  $eZ$ ,  $\nu W$ . The quoted limit assumes  $f = f' = \Lambda/m_{e^*}$ . See their Fig. 3 and Fig. 4 for the exclusion plots in the mass-coupling plane.
- 33 ABAZOV 08H search for single  $e^*$  production in  $p\bar{p}$  collisions with the decays  $e^* \rightarrow e\gamma$ . The  $e^*$  production is assumed to be described by an effective four-fermion interaction. See their Fig. 5 for the exclusion plot in the mass-coupling plane.
- 34 ACOSTA 05B search for single  $e^*$  production in  $p\bar{p}$  collisions with the decays  $e^* \rightarrow e\gamma$ .  $f = f' = \Lambda/m_{e^*}$  is assumed for the  $e^*$  coupling. See their Fig.3 for the exclusion limit in the mass-coupling plane.
- 35 ACHARD 03B result is from  $e^+e^-$  collisions at  $\sqrt{s} = 189\text{--}209$  GeV. See their Fig. 4 for the exclusion plot in the mass-coupling plane.
- 36 ABBIENDI 02G result is from  $e^+e^-$  collisions at  $\sqrt{s} = 183\text{--}209$  GeV.  $f = f' = \Lambda/m_{e^*}$  is assumed for  $e^*$  coupling. See their Fig. 4c for the exclusion limit in the mass-coupling plane.
- 37 CHEKANOV 02D search for single  $e^*$  production in  $ep$  collisions with the decays  $e^* \rightarrow e\gamma$ ,  $eZ$ ,  $\nu W$ .  $f = f' = \Lambda/m_{e^*}$  is assumed for the  $e^*$  coupling. See their Fig. 5a for the exclusion plot in the mass-coupling plane.

### Limits for Excited $e$ ( $e^*$ ) from $e^+e^- \rightarrow \gamma\gamma$

These limits are derived from indirect effects due to  $e^*$  exchange in the  $t$  channel and depend on transition magnetic coupling between  $e$  and  $e^*$ . All limits are for  $\lambda_\gamma = 1$ . All limits except ABE 89J and ACHARD 02D are for nonchiral coupling with  $\eta_L = \eta_R = 1$ . We choose the chiral coupling limit as the best limit and list it in the Summary Table.

For limits prior to 1987, see our 1992 edition (Physical Review **D45** S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;356</b>	95	38 ABDALLAH	04N DLPH	$\sqrt{s}= 161\text{--}208$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>310	95	ACHARD	02D L3	$\sqrt{s}= 192\text{--}209$ GeV

- 38 ABDALLAH 04N also obtain a limit on the excited electron mass with  $ee^*$  chiral coupling,  $m_{e^*} > 295$  GeV at 95% CL.

### Indirect Limits for Excited $e$ ( $e^*$ )

These limits make use of loop effects involving  $e^*$  and are therefore subject to theoretical uncertainty.

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
	39 DORENBOS...	89 CHRM	$\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e, \nu_\mu e \rightarrow \nu_\mu e$
	40 GRIFOLS	86 THEO	$\nu_\mu e \rightarrow \nu_\mu e$
	41 RENARD	82 THEO	$g-2$ of electron

- <sup>39</sup> DORENBOSCH 89 obtain the limit  $\lambda_\gamma^2 \Lambda_{\text{cut}}^2 / m_{e^*}^2 < 2.6$  (95% CL), where  $\Lambda_{\text{cut}}$  is the cutoff scale, based on the one-loop calculation by GRIFOLS 86. If one assumes that  $\Lambda_{\text{cut}} = 1$  TeV and  $\lambda_\gamma = 1$ , one obtains  $m_{e^*} > 620$  GeV. However, one generally expects  $\lambda_\gamma \approx m_{e^*} / \Lambda_{\text{cut}}$  in composite models.
- <sup>40</sup> GRIFOLS 86 uses  $\nu_\mu e \rightarrow \nu_\mu e$  and  $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$  data from CHARM Collaboration to derive mass limits which depend on the scale of compositeness.
- <sup>41</sup> RENARD 82 derived from  $g-2$  data limits on mass and couplings of  $e^*$  and  $\mu^*$ . See figures 2 and 3 of the paper.

## MASS LIMITS for Excited $\mu$ ( $\mu^*$ )

### Limits for Excited $\mu$ ( $\mu^*$ ) from Pair Production

These limits are obtained from  $e^+ e^- \rightarrow \mu^{*+} \mu^{*-}$  and thus rely only on the (electroweak) charge of  $\mu^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $\mu^*$  coupling is assumed to be of sequential type. All limits assume a dominant  $\mu^* \rightarrow \mu\gamma$  decay except the limits from  $\Gamma(Z)$ .

For limits prior to 1987, see our 1992 edition (Physical Review **D45** S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;103.2</b>	95	<sup>42</sup> ABBIENDI	02G OPAL	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type

• • • We do not use the following data for averages, fits, limits, etc. • • •

>102.8	95	<sup>43</sup> ACHARD	03B L3	$e^+ e^- \rightarrow \mu^* \mu^*$ Homodoublet type
--------	----	----------------------	--------	--

<sup>42</sup> From  $e^+ e^-$  collisions at  $\sqrt{s} = 183\text{--}209$  GeV.  $f = f'$  is assumed.

<sup>43</sup> From  $e^+ e^-$  collisions at  $\sqrt{s} = 189\text{--}209$  GeV.  $f = f'$  is assumed. ACHARD 03B also obtain limit for  $f = -f'$ :  $m_{\mu^*} > 96.6$  GeV.

### Limits for Excited $\mu$ ( $\mu^*$ ) from Single Production

These limits are from  $e^+ e^- \rightarrow \mu^* \mu$  and depend on transition magnetic coupling between  $\mu$  and  $\mu^*$ . All limits assume  $\mu^* \rightarrow \mu\gamma$  decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda\text{--}m_{\mu^*}$  plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45** S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;1090</b>	95	<sup>44</sup> CHATRCHYAN	11X CMS	$pp \rightarrow \mu\mu^* X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

	95	<sup>45</sup> ABAZOV	06E D0	$p\bar{p} \rightarrow \mu\mu^*$
> 221	95	<sup>46</sup> ABULENCIA,A	06B CDF	$p\bar{p} \rightarrow \mu\mu^*, \mu^* \rightarrow \mu\gamma$
> 180	95	<sup>47</sup> ACHARD	03B L3	$e^+ e^- \rightarrow \mu\mu^*$
> 190	95	<sup>48</sup> ABBIENDI	02G OPAL	$e^+ e^- \rightarrow \mu\mu^*$

- 44 CHATRCHYAN 11X search for single  $\mu^*$  production in  $pp$  collisions with the decay  $\mu^* \rightarrow \mu\gamma$ .  $f = f' = \Lambda/m_{\mu^*}$  is assumed. See their Fig. 2 for the exclusion plot in the mass-coupling plane.
- 45 ABAZOV 06E assume  $\mu\mu^*$  production via four-fermion contact interaction  $(4\pi/\Lambda^2)(\bar{q}_L\gamma^\mu q_L)(\bar{\mu}_L^*\gamma_\mu\mu)$ . The obtained limit is  $m_{\mu^*} > 618$  GeV ( $m_{\mu^*} > 688$  GeV) for  $\Lambda = 1$  TeV ( $\Lambda = m_{\mu^*}$ ).
- 46  $f = f' = \Lambda/m_{\mu^*}$  is assumed for the  $\mu^*$  coupling. See their Fig.4 for the exclusion limit in the mass-coupling plane. ABULENCIA,A 06B also obtain  $m_{\mu^*}$  limit in the contact interaction model with  $\Lambda = m_{\mu^*}$ ,  $m_{\mu^*} > 696$  GeV.
- 47 ACHARD 03B result is from  $e^+e^-$  collisions at  $\sqrt{s} = 189\text{--}209$  GeV.  $f = f' = \Lambda/m_{\mu^*}$  is assumed. See their Fig. 4 for the exclusion plot in the mass-coupling plane.
- 48 ABBIENDI 02G result is from  $e^+e^-$  collisions at  $\sqrt{s} = 183\text{--}209$  GeV.  $f = f' = \Lambda/m_{\mu^*}$  is assumed for  $\mu^*$  coupling. See their Fig. 4c for the exclusion limit in the mass-coupling plane.

### Indirect Limits for Excited $\mu$ ( $\mu^*$ )

These limits make use of loop effects involving  $\mu^*$  and are therefore subject to theoretical uncertainty.

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
• • •			We do not use the following data for averages, fits, limits, etc. • • •
	49 RENARD	82	THEO $g-2$ of muon
	49 RENARD	82	derived from $g-2$ data limits on mass and couplings of $e^*$ and $\mu^*$ . See figures 2 and 3 of the paper.

## MASS LIMITS for Excited $\tau$ ( $\tau^*$ )

### Limits for Excited $\tau$ ( $\tau^*$ ) from Pair Production

These limits are obtained from  $e^+e^- \rightarrow \tau^{*+}\tau^{*-}$  and thus rely only on the (electroweak) charge of  $\tau^*$ . Form factor effects are ignored unless noted. For the case of limits from  $Z$  decay, the  $\tau^*$  coupling is assumed to be of sequential type. All limits assume a dominant  $\tau^* \rightarrow \tau\gamma$  decay except the limits from  $\Gamma(Z)$ .

For limits prior to 1987, see our 1992 edition (Physical Review **D45** S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>103.2	95	50 ABBIENDI	02G OPAL	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
• • •				We do not use the following data for averages, fits, limits, etc. • • •
>102.8	95	51 ACHARD	03B L3	$e^+e^- \rightarrow \tau^*\tau^*$ Homodoublet type
		50		From $e^+e^-$ collisions at $\sqrt{s} = 183\text{--}209$ GeV. $f = f'$ is assumed.
		51		From $e^+e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = f'$ is assumed. ACHARD 03B also obtain limit for $f = -f'$ : $m_{\tau^*} > 96.6$ GeV.

## Limits for Excited $\tau$ ( $\tau^*$ ) from Single Production

These limits are from  $e^+e^- \rightarrow \tau^*\tau$  and depend on transition magnetic coupling between  $\tau$  and  $\tau^*$ . All limits assume  $\tau^* \rightarrow \tau\gamma$  decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling,  $\eta_L = \eta_R = 1$ . In most papers, the limit is expressed in the form of an excluded region in the  $\lambda-m_{\tau^*}$  plane. See the original papers.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;185</b>	95	<sup>52</sup> ABBIENDI	02G OPAL	$e^+e^- \rightarrow \tau\tau^*$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>180	95	<sup>53</sup> ACHARD	03B L3	$e^+e^- \rightarrow \tau\tau^*$
<sup>52</sup> ABBIENDI 02G result is from $e^+e^-$ collisions at $\sqrt{s} = 183\text{--}209$ GeV. $f = f' = \Lambda/m_{\tau^*}$ is assumed for $\tau^*$ coupling. See their Fig. 4c for the exclusion limit in the mass-coupling plane.				
<sup>53</sup> ACHARD 03B result is from $e^+e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = f' = \Lambda/m_{\tau^*}$ is assumed. See their Fig. 4 for the exclusion plot in the mass-coupling plane.				

## MASS LIMITS for Excited Neutrino ( $\nu^*$ )

### Limits for Excited $\nu$ ( $\nu^*$ ) from Pair Production

These limits are obtained from  $e^+e^- \rightarrow \nu^*\nu^*$  and thus rely only on the (electroweak) charge of  $\nu^*$ . Form factor effects are ignored unless noted. The  $\nu^*$  coupling is assumed to be of sequential type unless otherwise noted. All limits assume a dominant  $\nu^* \rightarrow \nu\gamma$  decay except the limits from  $\Gamma(Z)$ .

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;102.6</b>	95	<sup>54</sup> ACHARD	03B L3	$e^+e^- \rightarrow \nu^*\nu^*$ Homodoublet type
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		<sup>55</sup> ABBIENDI	04N OPAL	
<sup>54</sup> From $e^+e^-$ collisions at $\sqrt{s} = 189\text{--}209$ GeV. $f = -f'$ is assumed. ACHARD 03B also obtain limit for $f = f'$ : $m_{\nu_e^*} > 101.7$ GeV, $m_{\nu_\mu^*} > 101.8$ GeV, and $m_{\nu_\tau^*} > 92.9$ GeV. See their Fig. 4 for the exclusion plot in the mass-coupling plane.				
<sup>55</sup> From $e^+e^-$ collisions at $\sqrt{s} = 192\text{--}209$ GeV, ABBIENDI 04N obtain limit on $\sigma(e^+e^- \rightarrow \nu^*\nu^*) B^2(\nu^* \rightarrow \nu\gamma)$ . See their Fig.2. The limit ranges from 20 to 45fb for $m_{\nu^*} > 45$ GeV.				

### Limits for Excited $\nu$ ( $\nu^*$ ) from Single Production

These limits are from  $e^+e^- \rightarrow \nu\nu^*$ ,  $Z \rightarrow \nu\nu^*$ , or  $ep \rightarrow \nu^*X$  and depend on transition magnetic coupling between  $\nu/e$  and  $\nu^*$ . Assumptions about  $\nu^*$  decay mode are given in footnotes.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;213</b>	95	<sup>56</sup> AARON	08 H1	$ep \rightarrow \nu^*X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>190	95	<sup>57</sup> ACHARD	03B L3	$e^+e^- \rightarrow \nu\nu^*$
none 50–150	95	<sup>58</sup> ADLOFF	02 H1	$ep \rightarrow \nu^*X$
>158	95	<sup>59</sup> CHEKANOV	02D ZEUS	$ep \rightarrow \nu^*X$
>171	95	<sup>60</sup> ACCIARRI	01D L3	$e^+e^- \rightarrow \nu\nu^*$

- <sup>56</sup> AARON 08 search for single  $\nu^*$  production in  $ep$  collisions with the decays  $\nu^* \rightarrow \nu\gamma$ ,  $\nu Z$ ,  $eW$ . The quoted limit assumes  $f = -f' = \Lambda/m_{\nu^*}$ . See their Fig. 3 and Fig. 4 for the exclusion plots in the mass-coupling plane.
- <sup>57</sup> ACHARD 03B result is from  $e^+e^-$  collisions at  $\sqrt{s} = 189\text{--}209$  GeV. The quoted limit is for  $\nu_e^*$ .  $f = -f' = \Lambda/m_{\nu^*}$  is assumed. See their Fig. 4 for the exclusion plot in the mass-coupling plane.
- <sup>58</sup> ADLOFF 02 search for single  $\nu^*$  production in  $ep$  collisions with the decays  $\nu^* \rightarrow \nu\gamma$ ,  $\nu Z$ ,  $eW$ . The quoted limit assumes  $f = -f' = \Lambda/m_{\nu^*}$ . See their Fig. 1 for the exclusion plots in the mass-coupling plane.
- <sup>59</sup> CHEKANOV 02D search for single  $\nu^*$  production in  $ep$  collisions with the decays  $\nu^* \rightarrow \nu\gamma$ ,  $\nu Z$ ,  $eW$ .  $f = -f' = \Lambda/m_{\nu^*}$  is assumed for the  $e^*$  coupling. CHEKANOV 02D also obtain limit for  $f = f' = \Lambda/m_{\nu^*}$ :  $m_{\nu^*} > 135$  GeV. See their Fig. 5c and Fig. 5d for the exclusion plot in the mass-coupling plane.
- <sup>60</sup> ACCIARRI 01D search for  $\nu\nu^*$  production in  $e^+e^-$  collisions at  $\sqrt{s} = 192\text{--}202$  GeV with decays  $\nu^* \rightarrow \nu\gamma$ ,  $\nu^* \rightarrow eW$ .  $f = -f' = \Lambda/m_{\nu^*}$  is assumed for the  $\nu^*$  coupling. See their Fig. 4 for limits in the mass-coupling plane.

## MASS LIMITS for Excited $q$ ( $q^*$ )

### Limits for Excited $q$ ( $q^*$ ) from Pair Production

These limits are mostly obtained from  $e^+e^- \rightarrow q^*\bar{q}^*$  and thus rely only on the (electroweak) charge of the  $q^*$ . Form factor effects are ignored unless noted. Assumptions about the  $q^*$  decay are given in the comments and footnotes.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;338</b>	95	<sup>61</sup> AALTONEN 10H	CDF	$q^* \rightarrow tW^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		<sup>62</sup> BARATE 98U	ALEP	$Z \rightarrow q^*q^*$
> 45.6	95	<sup>63</sup> ADRIANI 93M	L3	$u$ or $d$ type, $Z \rightarrow q^*q^*$
> 41.7	95	<sup>64</sup> BARDADIN-...	92 RVUE	$u$ -type, $\Gamma(Z)$
> 44.7	95	<sup>64</sup> BARDADIN-...	92 RVUE	$d$ -type, $\Gamma(Z)$
> 40.6	95	<sup>65</sup> DECAMP 92	ALEP	$u$ -type, $\Gamma(Z)$
> 44.2	95	<sup>65</sup> DECAMP 92	ALEP	$d$ -type, $\Gamma(Z)$
> 45	95	<sup>66</sup> DECAMP 92	ALEP	$u$ or $d$ type, $Z \rightarrow q^*q^*$
> 45	95	<sup>65</sup> ABREU 91F	DLPH	$u$ -type, $\Gamma(Z)$
> 45	95	<sup>65</sup> ABREU 91F	DLPH	$d$ -type, $\Gamma(Z)$

<sup>61</sup> AALTONEN 10H obtain limits on the  $q^*q^*$  production cross section in  $p\bar{p}$  collisions. See their Fig. 3.

<sup>62</sup> BARATE 98U obtain limits on the form factor. See their Fig. 16 for limits in mass-form factor plane.

<sup>63</sup> ADRIANI 93M limit is valid for  $B(q^* \rightarrow qg) > 0.25$  (0.17) for up (down) type.

<sup>64</sup> BARDADIN-OTWINOWSKA 92 limit based on  $\Delta\Gamma(Z) < 36$  MeV.

<sup>65</sup> These limits are independent of decay modes.

<sup>66</sup> Limit is for  $B(q^* \rightarrow qg) + B(q^* \rightarrow q\gamma) = 1$ .

## Limits for Excited $q$ ( $q^*$ ) from Single Production

These limits are from  $e^+e^- \rightarrow q^*\bar{q}$ ,  $p\bar{p} \rightarrow q^*X$ , or  $pp \rightarrow q^*X$  and depend on transition magnetic couplings between  $q$  and  $q^*$ . Assumptions about  $q^*$  decay mode are given in the footnotes and comments.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;2490</b>	95	67 CHATRCHYAN11Y	CMS	$pp \rightarrow q^*X$ , $q^* \rightarrow qg$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 300–1260	95	68 ABAZOV	11F D0	$p\bar{p} \rightarrow q^*X$ , $q^* \rightarrow qZ, qW$
none 500–1580	95	69 AAD	10 ATLS	$pp \rightarrow q^*X$ , $q^* \rightarrow qg$
> 510	95	69 KHACHATRYAN...10	CMS	$pp \rightarrow q^*X$ , $q^* \rightarrow qg$
> 775	95	70 ABAZOV	06F D0	$p\bar{p} \rightarrow q^*X$ , $q^* \rightarrow qZ$
		71 ABAZOV	04C D0	$p\bar{p} \rightarrow q^*X$ , $q^* \rightarrow qg$

<sup>67</sup> CHATRCHYAN 11Y assume degenerate  $q^*$  with  $f_S = \Lambda/m_{q^*}$ .

<sup>68</sup> ABAZOV 11F search for vectorlike quarks decaying to  $W$ +jet and  $Z$ +jet in  $p\bar{p}$  collisions. See their Fig. 3 and Fig. 4 for the limits on  $\sigma \cdot B$ .

<sup>69</sup> AAD 10, KHACHATRYAN 10 search for heavy resonance decaying to 2 jets in  $pp$  collisions at  $\sqrt{s} = 7$  TeV.  $f_S = f = f' = 1$  is assumed.

<sup>70</sup> ABAZOV 06F assume  $q^*$  production via  $qg$  fusion and via contact interactions. The quoted limit is for  $\Lambda = m_{q^*}$ .

<sup>71</sup> ABAZOV 04C assume  $f_S = f = f' = \Lambda/m_{q^*}$ .

## MASS LIMITS for Color Sextet Quarks ( $q_6$ )

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;84</b>	95	72 ABE	89D CDF	$p\bar{p} \rightarrow q_6\bar{q}_6$

<sup>72</sup> ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color sextet quark is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. A limit of 121 GeV is obtained for a color decuplet.

## MASS LIMITS for Color Octet Charged Leptons ( $l_8$ )

$$\lambda \equiv m_{l_8}/\Lambda$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;86</b>	95	73 ABE	89D CDF	Stable $l_8$ : $p\bar{p} \rightarrow l_8\bar{l}_8$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>74</sup> ABT 93 H1  $e_8$ :  $ep \rightarrow e_8X$

<sup>73</sup> ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color octet lepton is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. The limit improves to 99 GeV if it always fragments into a unit-charged hadron.

<sup>74</sup> ABT 93 search for  $e_8$  production via  $e$ -gluon fusion in  $ep$  collisions with  $e_8 \rightarrow eg$ . See their Fig. 3 for exclusion plot in the  $m_{e_8}-\Lambda$  plane for  $m_{e_8} = 35-220$  GeV.

**MASS LIMITS for Color Octet Neutrinos ( $\nu_8$ )**

$$\lambda \equiv m_{\ell_8}/\Lambda$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;110</b>	90	<sup>75</sup> BARGER	89	RVUE $\nu_8: p\bar{p} \rightarrow \nu_8\bar{\nu}_8$

• • • We do not use the following data for averages, fits, limits, etc. • • •

none 3.8–29.8	95	<sup>76</sup> KIM	90	AMY $\nu_8: e^+e^- \rightarrow$ acoplanar jets
none 9–21.9	95	<sup>77</sup> BARTEL	87B	JADE $\nu_8: e^+e^- \rightarrow$ acoplanar jets

<sup>75</sup> BARGER 89 used ABE 89B limit for events with large missing transverse momentum. Two-body decay  $\nu_8 \rightarrow \nu g$  is assumed.

<sup>76</sup> KIM 90 is at  $E_{\text{cm}} = 50\text{--}60.8$  GeV. The same assumptions as in BARTEL 87B are used.

<sup>77</sup> BARTEL 87B is at  $E_{\text{cm}} = 46.3\text{--}46.78$  GeV. The limit assumes the  $\nu_8$  pair production cross section to be eight times larger than that of the corresponding heavy neutrino pair production. This assumption is not valid in general for the weak couplings, and the limit can be sensitive to its  $SU(2)_L \times U(1)_Y$  quantum numbers.

**MASS LIMITS for  $W_8$  (Color Octet  $W$  Boson)**

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>78</sup> ALBAJAR 89 UA1  $p\bar{p} \rightarrow W_8 X, W_8 \rightarrow Wg$

<sup>78</sup> ALBAJAR 89 give  $\sigma(W_8 \rightarrow W + \text{jet})/\sigma(W) < 0.019$  (90% CL) for  $m_{W_8} > 220$  GeV.

**REFERENCES FOR Searches for Quark and Lepton Compositeness**

AAD	11	PL B694 327	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	11E	PR D84 011101	G. Aad <i>et al.</i>	(ATLAS Collab.)
AARON	11C	PL B705 52	F. D. Aaron <i>et al.</i>	(H1 Collab.)
ABAZOV	11F	PRL 106 081801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	11	EPJ C71 1555	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
CHATRCHYAN	11X	PL B704 143	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	11Y	PL B704 123	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
KHACHATRYAN	11F	PRL 106 201804	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AAD	10	PRL 105 161801	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	10H	PRL 104 091801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
KHACHATRYAN	10	PRL 105 211801	V. Khachatryan <i>et al.</i>	(CMS Collab.)
Also		PRL 106 029902E	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRYAN	10A	PRL 105 262001	V. Khachatryan <i>et al.</i>	(CMS Collab.)
ABAZOV	09AE	PRL 103 191803	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	09	EPJ C60 1	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
AARON	08	PL B663 382	F.D. Aaron <i>et al.</i>	(H1 Collab.)
AARON	08A	PL B666 131	F.D. Aaron <i>et al.</i>	(H1 Collab.)
ABAZOV	08H	PR D77 091102R	V.M. Abazov <i>et al.</i>	(D0 Collab.)
SCHAEEL	07A	EPJ C49 411	S. Schael <i>et al.</i>	(ALEPH Collab.)
ABAZOV	06E	PR D73 111102R	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	06F	PR D74 011104R	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	06C	EPJ C45 589	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABULENCIA	06L	PRL 96 211801	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA,A	06B	PRL 97 191802	A. Abulencia <i>et al.</i>	(CDF Collab.)
ACOSTA	05B	PRL 94 101802	D. Acosta <i>et al.</i>	(CDF Collab.)
ABAZOV	04C	PR D69 111101R	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	04G	EPJ C33 173	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	04N	PL B602 167	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	04N	EPJ C37 405	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ACHARD	03B	PL B568 23	P. Achard <i>et al.</i>	(L3 Collab.)
BABICH	03	EPJ C29 103	A.A. Babich <i>et al.</i>	
ABBIENDI	02G	PL B544 57	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ACHARD	02D	PL B531 28	P. Achard <i>et al.</i>	(L3 Collab.)

ADLOFF	02	PL B525 9	C. Adloff <i>et al.</i>	(H1 Collab.)
CHEKANOV	02D	PL B549 32	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ACCIARRI	01D	PL B502 37	M. Acciarri <i>et al.</i>	(L3 Collab.)
AFFOLDER	01I	PRL 87 231803	T. Affolder <i>et al.</i>	(CDF Collab.)
BOURILKOV	01	PR D64 071701	D. Bourilkov	
CHEUNG	01B	PL B517 167	K. Cheung	
ABREU	00S	PL B485 45	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00P	PL B489 81	M. Acciarri <i>et al.</i>	(L3 Collab.)
AFFOLDER	00I	PR D62 012004	T. Affolder <i>et al.</i>	(CDF Collab.)
BARATE	98U	EPJ C4 571	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARGER	98E	PR D57 391	V. Barger <i>et al.</i>	
MCFARLAND	98	EPJ C1 509	K.S. McFarland <i>et al.</i>	(CCFR/NuTeV Collab.)
ABE	97T	PRL 79 2198	F. Abe <i>et al.</i>	(CDF Collab.)
DIAZCRUZ	94	PR D49 R2149	J.L. Diaz Cruz, O.A. Sampayo	(CINV)
ABT	93	NP B396 3	I. Abt <i>et al.</i>	(H1 Collab.)
ADRIANI	93M	PRPL 236 1	O. Adriani <i>et al.</i>	(L3 Collab.)
BARADIN...	92	ZPHY C55 163	M. Bardadin-Otwinowska	(CLER)
DECAMP	92	PRPL 216 253	D. Decamp <i>et al.</i>	(ALEPH Collab.)
PDG	92	PR D45 S1	K. Hikasa <i>et al.</i>	(KEK, LBL, BOST+)
ABREU	91F	NP B367 511	P. Abreu <i>et al.</i>	(DELPHI Collab.)
KIM	90	PL B240 243	G.N. Kim <i>et al.</i>	(AMY Collab.)
ABE	89B	PRL 62 1825	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	89D	PRL 63 1447	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	89J	ZPHY C45 175	K. Abe <i>et al.</i>	(VENUS Collab.)
ALBAJAR	89	ZPHY C44 15	C. Albajar <i>et al.</i>	(UA1 Collab.)
BARGER	89	PL B220 464	V. Barger <i>et al.</i>	(WISC, KEK)
DORENBOS...	89	ZPHY C41 567	J. Dorenbosch <i>et al.</i>	(CHARM Collab.)
BARTEL	87B	ZPHY C36 15	W. Bartel <i>et al.</i>	(JADE Collab.)
GRIFOLS	86	PL 168B 264	J.A. Grifols, S. Peris	(BARC)
JODIDIO	86	PR D34 1967	A. Jodidio <i>et al.</i>	(LBL, NWES, TRIU)
Also		PR D37 237 (erratum)	A. Jodidio <i>et al.</i>	(LBL, NWES, TRIU)
EICHTEN	84	RMP 56 579	E. Eichten <i>et al.</i>	(FNAL, LBL, OSU)
RENARD	82	PL 116B 264	F.M. Renard	(CERN)